



Patient Name: \_\_\_\_\_ Date: \_\_\_\_\_

### **COVID-19 Rapid IgG Antibodies test is: Negative/Positive**

This test detects IgG antibodies that develop in most patients within seven to 10 days after symptoms of COVID-19 begin. IgG antibodies remain in the blood after an infection has passed. These antibodies indicate that you may have had COVID-19 in the recent past and have developed antibodies that may protect you from future infection. It is unknown at this point how much protection antibodies might provide against reinfection.

### **COVID-19 Rapid IgM Antibodies test is: Negative/Positive**

This test detects IgM antibodies. IgM is usually the first antibody produced by the immune system when a virus attacks. A positive IgM test indicates that you may have been infected and that your immune system has started responding to the virus. When IgM is detected you may still be infected, or you may have recently recovered from a COVID-19 infection.

### **Disclaimer and important information:**

- This test has been authorized by the FDA for emergency use, pursuant to Section 564 of the Federal Food, Drug, and Cosmetic Act.
- Negative Results do not preclude acute SARS-CoV-2 infection. If acute infection is suspected, direct treating for SARS-CoV-2 is necessary.
- Results from antibody testing should not be used to diagnose or exclude acute SARS-CoV-2 infection.
- Positive results may be due to past or present infection with non-SARS-CoV-2 coronavirus strains, such as coronavirus HKU1, NL63, OC43, or 229E.
- This test is used for screening purposes only and not for diagnosing of COVID-19, for more definitive answer if you are currently infected and contagious, we highly recommend doing PCR testing by nasal swab for better virus detection.
- Manufacturer had applied for Emergency Use Authorization to the FDA.

### **Precautions and Recommendations:**

- You are still at risk for future infection if exposed to COVID-19 in the future.
- If your symptoms become more symptomatic, then seek medical help.
- Use universal precaution with mask and other protective equipment to reduce the chance of infection and spreading disease.
- Retesting may be indicated if symptoms develop.

Patient Acknowledgment: I understand that the IgG/IgM rapid test has limitations; included but not limited to there is a chance that you have had recent exposure in the last 7-14 days and that you did not develop antibodies yet. If that is the case; you may still have COVID-19 and still be contagious. In order to determine if someone is currently infected with COVID-19, a nasopharyngeal test (PCR) is recommended. The PCR testing is standard for identifying those with active COVID-19. You must speak to your employer for your company policy. We recommend compliance with the current CDC recommendations and this can be found at: [www.cdc.gov](http://www.cdc.gov)

Acknowledged by patient: \_\_\_\_\_

Acknowledged by Nurse: \_\_\_\_\_

Deland Wellness Center

122 S Amelia Ave Deland, FL 32724 (386)313.3200